

# Influence of Pentachlorophenol on Light Emission from Single Barnacle Muscle Fibers Preloaded with Aequorin

E. Edward Bittar and Jin-ru Wu

Department of Physiology, University of Wisconsin, Madison, WI 53706 USA

Interest has been aroused by studies on the pesticide pentachlorophenol (PCP), a known cytotoxicant (1,2), that showed PCB can stimulate the sodium efflux in barnacle muscle fibers at a concentration as low as 1  $\mu$ M and that its efficacy, which exceeds that of less-chlorinated phenols, depends on external pH ( $pH_e$ ). Additionally, the response of the sodium efflux to PCP is a sigmoidal function of external  $[Ca^{2+}]$ , the requirement for external  $Ca^{2+}$  being absolute (3). These findings have raised the question of whether PCP increases internal free  $Ca^{2+}$  by promoting the entry of  $Ca^{2+}$  from the outside into the myoplasm. This paper presents a study carried out with the photoprotein aequorin in an attempt to address the question of whether stimulation of the sodium efflux is due to a rise in internal free  $Ca^{2+}$  resulting from the entry of  $Ca^{2+}$  (4) and whether the efficacy of PCP depends on the number of chlorine atoms in its aromatic ring, as well as on the pH of the bathing medium ( $pH_e$ ).

Aequorin, rather than a  $Ca^{2+}$  dye substance such as Fura 2 or a  $Ca^{2+}$ -selective microelectrode, was chosen for several reasons. First, the aequorin method has been used with considerable success to study internal  $Ca^{2+}$  transients in skeletal muscle, most notably in barnacle fibers (5). Second, the aequorin signal is known to be relatively free from movement artifacts. Third, aequorin shows high sensitivity to  $Ca^{2+}$  and provides not only a sufficiently rapid response time but also a high signal-to-noise ratio. Fourth, aequorin is adequate for detecting internal  $Ca^{2+}$  transients over the physiological range of  $1 \times 10^{-7}$  to  $5 \times 10^{-6}$  M and lacks toxicity after injection (6).

We obtained specimens of the barnacle *Balanus nubilus* from Puget Sound, Seattle, Washington, and kept them in an aerated aquarium which was maintained at 10°–12°C throughout the study. The methods of dissection and cannulation of single fibers were essentially the same as those described by Bittar (7). The artificial sea water (ASW) used had the following composition: NaCl 465 mM; KCl 10 mM;  $MgCl_2$  10 mM;  $CaCl_2$  10 mM,  $NaHCO_3$  10 mM, and pH 7.8. We prepared solutions with varying concentrations of  $Ca^{2+}$  by raising or reducing NaCl in osmotically equivalent amounts. Solutions of PCP and other chlorinated phenols and phenol were

prepared by dissolving them in dimethyl sulfoxide (DMSO) before their addition to ASW. The final content of DMSO in ASW containing  $5 \times 10^{-4}$  M PCP was 0.5% (v/v). We used the buffers Hepes and Mops in lieu of  $HCO_3^-$  in experiments involving ASW at different pH values.

The microinjector used for loading the cannulated fibers with aequorin was of the type described by Bittar and Tallitsch (8). A solution of  $2\text{--}5 \times 10^{-4}$  M aequorin in a volume of 0.4–0.5  $\mu$ l was injected axially into these fibers with the aid of a micromanipulator and Palmer screw stand. Since the intrafiber fluid volume was on average 40  $\mu$ l, dilution by the myoplasm of the injected solution was approximately 100-fold. Aequorin was allowed to equilibrate for at least 1 hr. In those fibers poisoned with  $10^{-4}$  M ouabain, external application of this glycoside was carried out some 30 min before the application of pentachlorophenol (PCP) or any of its congeners.

The method used for recording aequorin light output was a modification of that described by Bittar and Keh (9) for monitoring firefly luciferase luminescence. Reading, display, and analysis of the picoammeter data was carried out with the aid of two software programs that were written in our laboratory by Patrick Hamilton. Experience with data processing revealed that analysis is best carried out by collating data based on experiments where basal light emission from the fibers is relatively the same. We estimated increments in light emission on the basis of maximal flash height; that is, by taking the difference between basal light emission and peak emission, expressed in nanoamperes. Significance levels were compared by using Student's *t*-test. A significance level of  $p < 0.05$  was selected. All experiments were carried out at an environmental temperature of 22°–24°C. All reagents used were analytical grade. Hepes, Mops, and ouabain were purchased from Sigma Chemical Company (St. Louis, Missouri). Pentachlorophenol and other chlorinated phenols and phenol were supplied by Aldrich Chemical Company (Milwaukee, Wisconsin).

In the first series of experiments with unpoisoned and ouabain-poisoned fibers, we investigated the effect on basal light output of external application of PCP at varying concentrations. Initially, it was necessary to determine whether the presence of DMSO (e.g., 0.5% in the bathing

Experiments show that the resting ouabain-insensitive sodium efflux in giant fibers from the barnacle *Balanus nubilus* is stimulated by external application of pentachlorophenol (PCP). This work has now been extended to include a study of muscle fibers preloaded with the  $Ca^{2+}$  indicator aequorin to determine whether PCP is able to increase light emission; and whether its potency depends on the number of chlorine atoms and external pH. The results obtained are as follows: 1) PCP causes a dose-dependent, multiphasic rise in light emission; the threshold concentration in fibers not poisoned with ouabain was in the low micromolar range. 2) The efficacy of PCP is considerably greater than that of less-chlorinated phenols and phenol. 3) The response to PCP is a sigmoidal function of external pH both in unpoisoned and ouabain-poisoned fibers. Reducing external pH potentiates its efficacy. 4) The response to PCP depends on the external  $Ca^{2+}$  concentration, and the requirement for  $Ca^{2+}$  is usually absolute. **Key words:** aequorin light emission, barnacle muscle fibers, pentachlorophenol. *Environ Health Perspect* 101:622–625(1993)

medium) influences basal light output from fibers (unpoisoned and ouabain poisoned) preloaded with aequorin. The results obtained in such experiments show that the presence of DMSO is without effect on basal light output. The characteristic signal observed in unpoisoned fibers to external application of PCP (e.g.,  $5 \times 10^{-4}$  M within 10 min) is an oscillatory rise in basal light emission, as illustrated in Figure 1a. A peak response to PCP is not immediate in onset. Such an experiment is shown in Figure 1b. Kinetics of this variety are also recorded with ouabain-poisoned fibers, but the onset of a peak response is more rapid and may be followed by a rather large, monophasic or almost monophasic  $Ca^{2+}$  transient, as shown in Figure 1c. Notice that the comparison drawn here is between fibers showing fairly similar baseline levels of light output.

It is noteworthy that the time to peak in unpoisoned fibers exposed to PCP shows a correlation with dosage only when the dose of PCP used falls in the high range. For example, time to peak in fibers exposed to  $10^{-4}$  M PCP averages  $183 \pm 73$  sec ( $n = 5$ ), a value that is not significantly


Address correspondence to E. E. Bittar, Department of Physiology, University of Wisconsin, 1300 University Avenue, Madison, WI 53706, USA.

Special thanks are due to O. Shimomura of Woods Hole Marine Laboratories, Woods Hole, Massachusetts, for supplying gifts of aequorin. This work was supported by NIH grant ESO5142. Received 1 March 1993; accepted 27 September 1993.

different from that obtained with fibers exposed to  $2.5$  or  $5 \times 10^{-4}$  M PCP, but significantly less than the values of  $392 \pm 27$  sec ( $n = 6$ ) and  $402 \pm 52$  sec ( $n = 6$ ) obtained with fibers exposed to  $7.5 \times 10^{-4}$  M and  $10^{-3}$  M PCP, respectively ( $p < 0.02$ ). On the other hand, an analysis of width at half-peak response fails to provide a similar correlation. Another feature that stands out is an increase in the frequency of internal transients in ouabain-poisoned fibers after exposure to PCP and that poisoned fibers are considerably more sensitive to PCP than unpoisoned fibers. For example, fibers poisoned with  $10^{-4}$  M ouabain and exposed to  $10^{-3}$  M PCP 30 min later show an increase in light emission, the magnitude of which averages  $224 \pm 29\%$  ( $n = 3$ ), whereas the increase observed in unpoisoned fibers exposed to  $10^{-3}$  M PCP averages  $76 \pm 17\%$  ( $n = 3$ ). The difference is significant.

In the second series, we tried to establish the concentration–response curve for the stimulatory effect of PCP on basal light emission using unpoisoned fibers. The time frame of the experiments was 10 min. Two features of the curve shown in Figure 2 are notable: first, light output is increased by PCP at a concentration as low as  $10 \mu\text{M}$ . This and a concentration of  $50 \mu\text{M}$  fall in the toxic range for aquatic organisms (10), as well as BF-2 cells derived from blue gill sunfish (11). Second, although a 10-fold elevation in PCP concentration fails to elicit a further increase in light output, a sharp inflexion of the curve occurs thereafter with a larger elevation in PCP concentration. Such results are consistent with the hypothesis that two mechanisms are operative: one involving a plasmalemma site which is saturated by PCP in low micromolar concentrations, and the other a myoplasmic site which is not readily saturated by PCP after its passage into the fiber interior.


In the third series of experiments, the response to PCP as a function of  $\text{pH}_e$  was examined. Since PCP is a weak acid with a  $\text{pK}_a$  of 4.8 (12), experiments were done to ascertain whether the efficacy of PCP depends in part on the pH of the suspending medium. These experiments involved the use of two different inert buffers, Hepes ( $\text{pK}_a$  7.5) and Mops ( $\text{pK}_a$  7.2). The results obtained with 10 mM Hepes ASW containing  $5 \times 10^{-4}$  M PCP show that light emission is a sigmoidal function of the medium pH. Light emission increases as  $\text{pH}_e$  is reduced. Similar results were obtained using unpoisoned and ouabain-poisoned fibers suspended in 10 mM Mops-ASW containing  $5 \times 10^{-4}$  M PCP. Summarized in Figure 3 are the results obtained with Mops as the buffer over a  $\text{pH}_e$  range of 8–6. Notice that  $\text{pH}_e$  (10 nM)




**Figure 1.** (A) Time-course of aequorin signals recorded from an unpoisoned fiber exposed to  $5 \times 10^{-4}$  M pentachlorophenol (PCP) at time indicated by arrow. Light output is expressed in nanoamperes. (B) A lag period precedes the onset of a marked outburst of light emission from aequorin. The PCP concentration was  $5 \times 10^{-4}$  M. (C) Time-course of a somewhat monophasic aequorin signal recorded from a fiber treated with  $10^{-4}$  M ouabain 30 min before exposure to  $5 \times 10^{-4}$  M pentachlorophenol at time indicated by arrow.

is fairly close to the threshold value for ouabain-poisoned fibers. Also notice that the midpoint of the slope is about 100 nM  $\text{H}^+$ . In a typical experiment, an unpoisoned fiber suspended in 10 mM Mops-ASW at pH 7 (i.e., 100 nM) shows a multiphasic rise in light emission when exposed to  $5 \times 10^{-4}$  M PCP, as illustrated in Figure 4a. The rise is from approximately 15 nA to less than 100 nA over less than 10 min. Multiphasic behavior is also seen in ouabain-poisoned fibers suspended in 10 mM Mops containing  $5 \times 10^{-4}$  M PCP at pH 6 (1000 nM), but the peak response is larger, as illustrated in Figure 4b. Notice that the basal level of light output before the addition of PCP is practically the same as that of the fiber in the preceding experiment.

Next, we studied the response to PCP as a function of chlorine atoms. The potency of chlorinated phenols is known




**Figure 2.** Concentration–response curve for the stimulatory effect of pentachlorophenol on light emission from aequorin in unpoisoned fibers. Vertical bars indicate  $\pm 1$  SEM; the number of measurements carried out is in parentheses. All fibers used were isolated from the same barnacle specimen (log–log plot).



**Figure 3.** (A) The dependency on  $\text{pH}_e$  of the magnitude of the increase in light emission in unpoisoned fibers and (B) ouabain-poisoned fibers in response to external application of  $5 \times 10^{-4}$  M pentachlorophenol in 10 mM Mops-ASW. Each plotted point in panel A is the mean of six measurements, except that at pH 6. Vertical bars indicate  $\pm 1$  SEM. The fibers used were isolated from three barnacle specimens taken from the same batch (semilog plot).

to increase with chlorination of the parent phenol (3,10,13). Figure 5 shows a representative experiment carried out with phenol. External application of  $5 \times 10^{-4}$  M phenol in ASW containing 10 mM Hepes at pH 7.4 produces a multiphasic rise in






**Figure 4.** Time-course of a multiphasic aequorin response of an ouabain-poisoned fiber suspended in 10 mM Mops-ASW at (A) pH 7 and (B) pH 6 to external application of  $5 \times 10^{-4}$  M pentachlorophenol (PCP) at time indicated by arrow. Notice the brief lag period preceding the onset of a rise in basal light emission in panel A.

light emission, but the maximum magnitude of this response is rather small. In view of such results, experiments were done in which a fixed concentration of  $5 \times 10^{-4}$  M of phenol, dichlorophenol, trichlorophenol, tetrachlorophenol, and pentachlorophenol was used for purposes of comparing their efficacy. Shown in Figure 6 is clear-cut evidence that an increase in light output in response to the application of these substances in equimolar concentration is a function of the number of chlorine atoms, pentachlorophenol being the most efficacious. To be certain of this conclusion, the experiments were repeated. The results obtained confirmed our conclusion.

Finally, experiments were carried out to determine whether the response to PCP depends on external  $\text{Ca}^{2+}$ . Using the recent results that the stimulatory response of the sodium efflux to PCP is a sigmoidal function of external  $\text{Ca}^{2+}$  and that the requirement for external  $\text{Ca}^{2+}$  is absolute (3), studies with ASW containing  $\text{Ca}^{2+}$  at varying concentrations were undertaken. Figure 7 illustrates that the magnitude of the observed rise in light emission after the application of  $5 \times 10^{-4}$  M PCP depends on the external  $\text{Ca}^{2+}$  concentration and that PCP is ineffective in the nominal absence of  $\text{Ca}^{2+}$  in the bathing medium. Other experiments, however, indicate that, on occasion, a small increase in basal light emission is observed with  $5 \times 10^{-4}$  M PCP. This is accounted for by assuming that some  $\text{Ca}^{2+}$  is retained in the fluid of the




**Figure 5.** Time-course of the multiphasic response obtained by external application (at time indicated by arrow) of  $5 \times 10^{-4}$  M phenol. Notice that the outbursts in light emission are rather small.


channels of the transverse tubular system of these fibers.

The experiments described here show quite clearly that PCP is able to increase light emission from fibers preloaded with aequorin and that the threshold concentration of PCP is practically the same as that required for stimulation of the basal sodium efflux (3). The failure of basal light emission to rise after the addition of PCP to nominally  $\text{Ca}^{2+}$ -free ASW is in agreement with the results of Nwoga and Bittar (3), who found that the sodium efflux rises after exposure of these fibers to PCP only if external  $\text{Ca}^{2+}$  is present in the bathing medium. Although the provisional conclusion which emerges is that "trigger"  $\text{Ca}^{2+}$  derives from the bathing medium, the possibility still remains that a fraction of the  $\text{Ca}^{2+}$  comes from internal storage sites, such as the sarcoplasmic reticulum. This possibility is not unlikely considering evidence that ryanodine and TMB, both of which act as blockers of the sarcoplasmic reticulum  $\text{Ca}^{2+}$  release channel, are able to reduce the response of the sodium efflux to PCP (3). It could be that the increase in internal free  $\text{Ca}^{2+}$  caused by PCP is partly the result of  $\text{IP}_3$ -promoted release of  $\text{Ca}^{2+}$  from the sarcoplasmic reticulum, but there is evidence that the ouabain-insensitive sodium efflux is unresponsive to the injection of  $\text{IP}_3$  (14). This is also true of thapsigargin (Bittar EE, unpublished data). Another possibility is that PCP, a known uncoupler (15,16), might cause the release of  $\text{Ca}^{2+}$  from mitochondria, as well as a reduction in internal ATP-Mg. However, neither possibility seems likely, partly because the classical uncoupler 2,4-dinitrophenol is without effect on the sodium efflux (Bittar EE, unpublished data), and partly because phosphagen occurs in abundance in these fibers [e.g.,  $^{31}\text{P}$ -NMR reveals an ArP content of 24 mmol/kg fiber water (17)].

There is circumstantial evidence favoring the view that the  $\text{Ca}^{2+}$  channel (L-type in barnacle fibers) is a primary point of action of PCP and that external  $\text{Ca}^{2+}$



**Figure 6.** Comparison of the potency of phenol and four chlorinated derivatives. The ouabain-poisoned fibers were suspended in 10 mM Hepes-ASW at pH 7.4. The concentrations used were equimolar ( $5 \times 10^{-4}$  M). Vertical bars indicate  $\pm 1$  SEM; the number of individual experiments carried out is in parentheses. The fibers used in these experimental sessions were isolated from three barnacle specimens taken from the same batch.



**Figure 7.** The dependency of light emission from aequorin in ouabain-poisoned fibers treated with  $10^{-3}$  M pentachlorophenol externally on the external  $\text{Ca}^{2+}$  concentration. Vertical bars indicate  $\pm 1$  SEM; the number of measurements recorded is in parentheses. The fibers used were isolated from the same barnacle specimen.

reaches the myoplasm via this pathway (3). However, this cannot be the complete picture: for example, to account more fully for the  $\text{Ca}^{2+}$  transients recorded with aequorin, one has to explain the part played by the  $\text{Na}^+-\text{Ca}^{2+}$  exchanger in maintaining cellular  $\text{Ca}^{2+}$  homeostasis in the face of an increased  $\text{Ca}^{2+}$  influx and a reduced sodium gradient in ouabain-poisoned fibers. This is precisely the rationale for using  $10^{-4}$  M ouabain; that is, to maximally inhibit the membrane  $\text{Na}^+-\text{K}^+$ -ATPase system, thereby raising the internal sodium concentration in these fibers (18). Thus, for example, under conditions where  $\Delta \tilde{\mu}_{\text{Ca}^{2+}} > 3\Delta \tilde{\mu}_{\text{Na}^+}$ , the energy provided by the  $\text{Ca}^{2+}$  electrochemical gradient would be expected to drive the sodium efflux (19). In other words, the marked outbursts of  $\text{Ca}^{2+}$  transients observed, particularly in ouabain-poisoned fibers, is

largely related to the operation of the  $\text{Na}^+-\text{Ca}^{2+}$  exchanger in the reverse mode. Because ATP behaves as a positive effector of this exchanger in barnacle fibers (20), we suggest that PCP stimulates reverse  $\text{Na}^+-\text{Ca}^{2+}$  exchange as the result of transiently raising the internal ATP level. This is likely to be the case if PCP acts as a competitive inhibitor of certain ATP-dependent processes. Evidence supporting this possibility comes from recent studies carried out with the firefly luciferase reaction *in vitro* (Xiang Z, Bittar EE, unpublished data). PCP is found to increase the  $K_m$  for ATP but not affect  $V_{\max}$ . Because a specific and powerful inhibitor of the  $\text{Na}^+-\text{Ca}^{2+}$  exchanger is not yet available (21), this rather interesting problem cannot be addressed in a direct manner at this time.

The pH-reaction profile is a sigmoid curve with a mid-point of about pH 7. This is compatible with a mechanism involving ionization of a group located in an active site, e.g., the  $\text{Ca}^{2+}$ -channel protein that facilitates the passage of external calcium into the myoplasm after interaction of PCP at this active site. An additional explanation of the observed dependency of PCP potency on  $\text{pH}_e$  is to invoke the law of nonionic diffusion (22,23). In other words, protonation of PCP renders it more lipophilic, thus leading to greater entry of the neutral species (e.g., into the plasmalemma compartment *per se*). This could bring about changes in lipid bilayer organization. Thus, to understand more fully the mechanism underlying the action of PCP, information is required not only about the role of the  $\text{Ca}^{2+}$  channel but also about whether PCP affects membrane fluidity. Until now, the few studies carried out with synthetic lipid bilayer membranes and biological membranes indicated that PCP in the micromolar range alters the physical properties of these systems (24–26). Another approach is to use an equilibrium model: this shows that at pH values less than pH 7 the distribution of the weak acid, e.g., between octanol and water, is dominated by the neutral species and does not depend on ionic strength (27). Such studies remain meager but are in keeping with the classical studies of Goodnight, who was the first to demonstrate that PCP can be toxic and that lowering pH potentiates this toxicity (28).

Finally, this study draws attention to the possibility that the cytotoxic action of PCP may be chiefly due to a raised cytosolic free  $\text{Ca}^{2+}$  concentration (29). Although human deaths have taken place after acute industrial, accidental, or suicidal exposure to PCP, the minimal lethal blood concen-

tration remains unknown (2). Whole-body exposure to PCP and urinary levels in the general population are very low [0.001–0.17 mg/person/day and 1–10  $\mu\text{g/l}$ , respectively (30)]. Aquatic species show extreme susceptibility to PCP toxicity. This is borne out by evidence of an  $\text{LC}_{50}$  value for PCP of about 1  $\mu\text{M}$  (31,32), which is a concentration close to that found to cause a rise in internal free  $\text{Ca}^{2+}$  in barnacle fibers.

## REFERENCES

- Rhao KR, ed. Pentachlorophenol: chemistry, pharmacology and environmental toxicology. New York:Plenum Press, 1978.
- Exon JH. A review of chlorinated phenols. *Vet Hum Toxicol* 26:508–520(1984).
- Nwoga J, Bittar EE. An investigation of the sensitivity of the ouabain-insensitive sodium efflux in single barnacle muscle fibers to pentachlorophenol. *Toxic Appl Pharmacol* 108:330–341(1991).
- Wu JR, Zhou Z, Bittar EE. Abolition with chloramine-T of inactivation in barnacle muscle fibres results in stimulation of the ouabain-insensitive sodium efflux. *Biochim Biophys Acta* 1112:99–104(1992).
- Ashley CC, Ridgway EB. On the relationship between membrane potential, calcium transient and tension in single barnacle muscle fibres. *J Physiol* 209:105–130(1970).
- Campbell AK. Chemiluminescence. Chichester, England:Ellis Horwood Ltd., 1988;384.
- Bittar EE. The barnacle muscle fibre as a model system for the investigation of the ouabain-insensitive sodium efflux and hormonal actions. *Progr Neurobiol* 20:1–54(1983).
- Bittar EE, Tallitsch RB. Stimulation by aldosterone of the sodium efflux in barnacle muscle fibres. *J Physiol* 250:337–346(1975).
- Bittar EE, Keh T. An investigation of myoplasmic magnesium adenosine triphosphate in barnacle muscle fibres with the firefly method. *J Physiol* 302:73–88(1980).
- Kozak VP, Simsiman GV, Chesters G, Stensby D, Harkins J. Reviews of the environmental effects of pollutants. XI. Chlorophenols. Cincinnati, OH:U.S. Environmental Protection Agency, 1979.
- Babich H, Borenfreund E. *In vitro* cytotoxicity of organic pollutants to bluegill sunfish (BF-2) cells. *Environ Res* 42:229–237(1987).
- Kaiser KLE, Valdmann I. Apparent octanol/water partition coefficients of pentachlorophenol as a function of pH. *Can J Chem* 60:2104–2106(1982).
- Ahlborg UG, Thunberg T. Chlorinated phenols: occurrence, toxicity, metabolism and environmental impact. *CRC Crit Rev Toxicol* 7:1–35(1980).
- Bittar EE, Huang YP. Lack of effect on the sodium efflux of the microinjection of D-Ins (1,4,5)  $\text{P}_3$  into ouabain-poisoned barnacle muscle fibres. *Biochim Biophys Acta* 1106:297–302(1992).
- Weinbach EE. The effect of pentachlorophenol on oxidative phosphorylation. *J Biol Chem* 210:545–550(1954).
- Weinbach EC, Garbus J. The interaction of uncoupling phenols with mitochondria and with mitochondrial protein. *J Biol Chem* 240:1811–1819(1965).
- Hansen J, Sharpe T, Bittar EE. Phosphate metabolites in single barnacle muscle fibers investigated by phosphorus-31 nuclear magnetic resonance. *Comp Biochem Physiol* 83B:875–879(1986).
- White JF, Hinke JAM. Use of the sodium microelectrode to define sodium efflux and the behavior of the sodium pump in the frog sartorius. In: *Ion and enzyme electrodes in toxicology and medicine* (Kessler M, Clark LC, Lubbes DW, Silver IA, Simon W, eds). Baltimore, MD:University Park Press, 1976;355–363.
- Allen TJ, Noble D, Reuter H, eds. Sodium-calcium exchange. Oxford:Oxford University Press, 1989.
- Bittar EE, Huang YP. ATP as a positive effector of the sodium efflux in single barnacle muscle fibers. *Biochim Biophys Acta* 1070:332–342(1991).
- Philipson KD. The cardiac  $\text{Na}^+-\text{Ca}^{2+}$  exchanger. In: *Calcium and the heart* (Langer G, ed). New York:Raven Press, 1990;85–108.
- Kempton RT. The differences in the elimination of neutral red and phenol red by the frog kidney. *J Cell Comp Physiol* 14:73–81(1939).
- Jacobs MH. Some aspects of cell permeability to weak electrolytes. *Cold Spring Harbor Symp Quant Biol* 8:30–39(1940).
- Danner J, Resnick H. Use of the fluorescent probe 1-anilino-8-naphthalene sulfonate to monitor the interactions of chlorophenols with phospholipid membranes (liposomes). *Biochem Pharmacol* 29:2471–2475(1980).
- Duxbury CL, Thompson JE. Pentachlorophenol alters the molecular organization of membranes in mammalian cells. *Arch Environ Contam Toxicol* 16:367–373(1987).
- Smejtek P, Barstad W, Wang S. Pentachlorophenol-induced change of  $\beta$ -potential and gel-to-fluid transition temperature in model lecithin membranes. *Chem-Biol Interact* 71:37–61(1989).
- Westall JC, Leuenberger C, Schwartzbach RP. Influence of pH and ionic strength in the aqueous-nonaqueous distribution of chlorinated phenols. *Environ Sci Technol* 19:193–198(1985).
- Goodnight CJ. Toxicity of sodium pentachlorophenolate and pentachlorophenol to fish. *Ind Eng Chem* 34:868–872(1942).
- Orrenius SS, McConkey DJ, Bellomo G, Nicotera P. Role of the  $\text{Ca}^{2+}$  in toxic cell killing. *Trends Pharmacol Sci* 10:281–285(1989).
- U.S. EPA. Ambient water quality criteria for pentachlorophenol. EPA 440-5-80-065. Washington, DC:U.S. Environmental Protection Agency, 1980.
- Cardwell RD, Foreman DG, Payne TR, Wilbur DJ. Acute toxicity of selected toxicants to six species of fish. EPA-600/3-76-008. Washington, DC:U.S. Environmental Protection Agency, 1976;1–117.
- Pierce RH. Fate and impact of pentachlorophenol in a fresh water ecosystem. EPA-600/3-78-063. Washington, DC:U.S. Environmental Protection Agency 1978;1–61.